

---

**A STUDY ON ABUNDANCE AND GROUP DIVERSITY OF SOIL MICROARTHROPODS AT  
FOUR DIFFERENT SOIL HABITATS IN NORTH DINAJPUR, WEST BENGAL INDIA**

S. K. Sarkar<sup>1</sup>, K. Chakrobarty<sup>2</sup> and M. N. Moitra<sup>3\*</sup>

<sup>1</sup>Pundari High School (H.S.), Dakshin Dinajpur, West Bengal, India; <sup>2</sup>Department of Zoology, Faculty of Science, Gour Banga University, West Bengal, India; <sup>3</sup>Department of Zoology, P. D. Women's College, West Bengal, India.

\*Corresponding author

**Abstract**

Sampling was conducted at four different sites i.e., an agricultural field, a river basin, a brick field and a forest floor from the district of Uttar Dinajpur, West Bengal, India. Though abundance was higher at the forest site, the group diversities during summer, monsoon and post-monsoon seasons were highest at the river basin. Site-specific and temporal variation of abundance was significant among the sites ( $p < 0.05$ ).

**Key words:** soil micro-arthropods, abundance, group diversity, agricultural field, forest.

**Introduction**

Soil micro-arthropods play significant role in ecological dynamics in soil and therefore have drawn attention of workers around the globe including India. (Crossley and Coleman, 1999; Colman et al., 2004; Devi and Singh, 2006; Chitrapati and Singh, 2006). Rutigliano et al., (2013) conducted works on the impact of fire on microarthropods. Impact of earthworm movements on microarthropods was investigated by Cameron et al., (2013). Bokhorst et al., (2014) studied the impact of mosses and shrubs on the soil dwellers in a boreal forest. Callejas et al., (2015) investigated the relationship between the soil microarthropods and the higher trophic level. The bio-indicator value of micro-arthropods in a forest ecosystem was assessed by Calugar and Ivan (2016) while, Lakshmi and Joseph (2016) worked on the utility of microarthropods as indicators of soil quality in a tropical garden.

The present study was taken up to examine the seasonal variation of microarthropod groups which are abundant in the alluvial plains of a part of northern parts of Bengal and also to assess the variability in group diversity in the region. The work is expected to provide a data base for the assessment of soil quality in the selected region.

**Material and Methods**

Soil samples were collected from four different sites during March, 2015 to February, 2016, with an interval of 30 days. Fifteen cores (5cm diameter) samples were collected from each of the five sub-plots of 1 m<sup>2</sup> area were selected at each site. Extraction was run using tullgren funnel apparatus as modified by Macfadyen (1953). Microarthropod groups were sorted from the extract and preserved in 80% alcohol. Mean values of four collection effort

per season (per 15 cores) were taken for comparison.

Shannon's Index following Cancela da Fonseca and Sarkar (1998) was used to work out group diversity. Logarithmic transformations of data, as per need of parametric statistics were made accordingly.

### **Collection sites**

A description of the sites has been given in an earlier publication (Sarkar et al., 2014). The sites are: S1. an agricultural field; S2. a brick field S3. a river basin region; S4. a forest area. All the sites are situated in the District Uttar Dinajpur, West Bengal, India.

### **Results and Discussion**

The observation is based upon a total collection of 720 cores (5cm diameterX5cm depth) from four sites during one year period. Seasonal abundance for each site was estimated taking a mean of total collection of three month for each season (individuals in 15 cores/season at a site). For analyses, major groups were divided as: 1.Oribtida, 2. Other mites, 3. Collembola, and 4. Other microarthropods (Cancela da Fonseca and Sarkar, 1998). Other mites included orders like mesostigmata, prostigmata and astigmata. Aranae and other insect orders like hymenoptera (mostly ants), diptera, coleoptera, (larval forms + nymphs mostly), protura, diplura etc excluding collembolan were grouped into other microarthropods.

One way ANOVA showed no statistically significant difference among the sites in variance of abundance while the mean abundances also did not differ significantly as the Tukey test indicated (Tables 1, 2). The variation however appeared significant ( $p < 0.05$ ) when sites and seasons were considered together in two-way ANOVA (Table 3). The numerical abundances of microarthropods were relatively higher during the post monsoon and lower during the summer at all the sampling sites (Figure 1). Bhattacharya

and Ray choudhury (1979), Bhattacharya et al., (1980) reported similar findings in their studies.

Oribatida was the most abundant order of microarthropods followed by collembolans in overall collection and in most of the collection efforts. Relative abundances of oribatid mites varied from 38.1% (S4) to 28.52% (S3). Relative abundance of other mites ranged from 14.47% to 22.63% while, the same of collembolans varied from 22.18% to 26.27% (Figure 2). Though the abundance of meso-stigmatid was high, the other two groups were very few in numbers and found to be absent in most collections efforts. This observation matches with the observations made by Sanyal (1982), Bhattacharya and Chakraborti (1994), Ghosh and Roy (2004), Joy (2006) and Moitra et al., (2007, 2013).

Group diversity was estimated upon seasonal collection. In summer, monsoon and post-monsoon, it was highest at S3 which is located at a river basin. More or less consistent availability of moisture might have rendered favourable effects on all the groups of microarthropods taken into account here in comparison to other sites. Higher group diversity at the sides of a sewage canal when compared to a wasteland, a roadside and a forest was reported by Moitra et al., (2013). Though the highest abundance of microarthropods was recorded as is expected for a forest site, group diversity was comparatively lower here. It may be due to the current procedure of calculation diversity where variety is fixed and it is rather evenness that would contribute more in the value. The observation therefore, probably indicates the greater dominance of oribatids as a major community of soil mesofauna in comparison to other sites sampled.

### **Conclusion**

In the current study, overall site-wise variation appeared minimal, but, seasons exhibited a significant variable effects on the abundances of






Figure 2. Relative abundance (%) of major microarthropod groups at the sampling sites.




Figure 3. Seasonal variation of group diversity at the sampling sites.

micro-arthropod communities when combined with plots. The efficiencies of various microarthropod extraction procedures are highly variable (Wallwork, 1976; Edwards, 1991; Akoijam, 2014) and several factors may have complex inter-related impacts on abundance, diversity and distribution of microarthropods (Cameron et al., 2013; Bokhorst et al., 2014). Further study therefore appears necessary to

have a better understanding of the edapho-ecological characteristics of the area selected.

#### **Acknowledgements**

Authors would like to express their sincere gratitude to all the staff of Dept. of Zoology, Gour Banga University for their cooperation throughout the current work.

#### **References**

- Akoijam, R., Bhattacharyya, B., and Mrangmei, L. (2014). Tullgren funnel – an efficient device for extracting soil microarthropods. *Environment and Ecology*. **32(2)**: 474-476.
- Bhattacharya, T., Bhattacharya, J., and Banerjee, R. (1980). A preliminary survey of the effect of smoke emission from a husking mill on the soil microarthropods. Newsletter, *Soil Biology & Ecology*. **1(2)**: 15-17.
- Bhattacharya, T. and Chakraborti, P. (1994). Community structure of soil Oribatida of a young Rubber plantation and an adjacent waste land in Tripura (India). In : *Advances in Ecology and Environmental Science*. PC Mishra, N Behera, BK Senapati, BC Guru (eds). pp. 65-77.
- Bhattacharya T. and Raychoudhuri D N. (1979). Monthly variation in the density of soil microarthropods in relation to some climatic and edaphic factors. *Entomon*. **4(4)**: 313-18.
- Bokhorst, S., Wardle, D.A., Nilsson, M.C. and Gundale, M.J. (2014). Impact of understory mosses and dwarf shrubs on soil microarthropods in a boreal forest Chronosequence. *Plant and Soil*. **379(1-2)**: 121-133.
- Călugăr, A. and Ivan, O. (2016). Soil microarthropods and their bioindicator value regarding the bio-edaphic conditions in forest ecosystems of Danube delta. *Seria Științele Vieții*. **26(2)**: 215-219.
- Callejas-Chavero, A., Castaño-Meneses, G., Razo-González, M., Pérez-Velázquez, D., Palacios-Vargas, J.G. and Flores-Martínez, A. (2015). *Journal of insect science*. **15(1)**: 1-9.
- Cameron E.K., Proctor, H.C. and Bayne, E.M. (2013). Effects of an Ecosystem Engineer on Belowground Movement of Microarthropods. *PLoS ONE* **8(4)**: e62796. doi:10.1371/journal.pone.0062796
- Cancela da Fonseca, J.P. and Sarkar, S. (1998). Soil microarthropods in two different managed ecological systems (Tripura, India). *Applied Soil Ecology*. **9**: 105-107.
- Chitrapati, C. and Singh, T.B. (2006). The role of abiotic factors in the distributional patterns of acarina and collembola in the sub-tropical forest ecosystem of Manipur. *Indian. J. Environ. & Ecoplan*. **12(1)**: 39-45.
- Coleman, D.C. and Crossley, D.A. (Jr) and Hendrix, P.F. (2004). *Fundamentals of soil ecology*. Elsevier Academic Press, New York.
- Crossley, D.A.(Jr) and Coleman, D.C. (1999). Microarthropods. In: *Handbook of Soil Science*. ME Sumner (ed). CRC Press, Boca Raton. pp. C-59 - C-65.

- Devi, K.L., and Singh, T.B. (2006). Population fluctuation of soil mites in relation to some important abiotic factors in the populations of soil-inhabiting invertebrates. *Agriculture, Ecosystems and Environment*. **34**: 145–176.
- Ghosh, T.C. and Roy, S. (2004). Distribution and diversity of acarina community three tea gardens at different altitudes of Darjeeling Himalayas. *Proc. Zool. Soc. Calcutta*. **57**: 87-93.
- Lakshmi, G. and Joseph, A. (2016). Soil microarthropods as indicators of soil quality of tropical home gardens in a village in Kerala, India. *Agro-forestry Systems*. pp 1-12. DOI 10.1007/s10457-016-9941-z
- MacFadyen, A. (1953). Notes on methods for the extraction of small soil arthropods. *J. Animal Ecol.* **22**: 65- 77.
- Moitra, M.N., Sanyal, A.K. and Chakrabarti, S. (2007). Variation of group diversity in soil microarthropod community at different altitudes in the Darjeeling Himalayas, West Bengal, India. *J. Environ & Sociobiol.* **4(2)**: 163-168.
- Moitra, M.N., Banerjee, S. and Sanyal, A.K. (2013). A study on group diversity as an assessment of biodiversity in soil microarthropod communities at four different habitats in and around Kolkata, pine forest ecosystem in Manipur, N. E. India. *J. Curr. Sci.* **9(2)**: 673 – 678.
- Edwards, C.A., (1991). The assessment of West Bengal, India. In: *Biotechnology for people*, Ed. S. Mukherjee. pp. 30-38.
- Rutigliano, F.A., Migliorini, M., Maggi, O., Ascoli, D., Fanciulli, P. P. and Persiani, A.M. (2013). Dynamics of fungi and fungivorous microarthropods in aMediterranean maquis soil affected by experimental fire. *European Journal of Soil Biology.* **56**: 33-43.
- Sanyal, A.K. (1982). Soil oribatid mites and their relation with soil factors in West Bengal. *J. Soil Biol, Ecol.* **2(1)**: 8-17.
- Sarkar, S.K., Chakraborty, K. and Moitra, M.N. (2014). Observation on abundance and group diversity of soil microarthropods with special reference to acarines at four differently used soil habitats. *International Journal of Scientific and Research Publications.* **4(11)**: 1-12.
- Wallwork, J.A. (1976). *The distribution and diversity of soil fauna*. Acad. Press Inc. Ltd. London. pp. 1-355.